

# Hepatitis B Immune Globulin

**Vaccines have saved more lives in Canada in the last 60 years than any other medical intervention. Vaccines help your immune system to recognize and fight bacteria and viruses that cause diseases.**

## What is hepatitis B immune globulin?

Hepatitis B (HB) immune globulin has large amounts of HB antibodies taken from donated human blood. Antibodies are proteins that a person's immune system makes to fight viruses or bacteria.

HB immune globulin provides immediate, short-term protection against the HB virus. It can prevent the disease or make the disease less severe. HB immune globulin works best if given:

- Within 48 hours after exposure to the HB virus; or
- Within 14 days of sexual contact with an infected individual.

## Is HB immune globulin safe?

Yes, HB immune globulin is a very safe human blood product. Canadian Blood Services screens and tests every blood donation. Since blood screening and testing began, there have been no reports of blood-borne infections like HB in persons receiving HB immune globulin.

## Who should get HB immune globulin?

If you or your child have potentially been exposed to the HB virus, and have not received the HB vaccine in the past, you may be advised to get HB immune globulin especially for:

- People who had unprotected sex with a person who has HB infection.
- Sexual assault victims.
- Infants whose mothers have HB.
- Infants whose mothers are at high risk of infection with HB, such as intravenous drug users or sex trade workers.

A dose of HB vaccine may be given at the same time as HB immune globulin. More doses of HB vaccine may be given later to provide full, long-term protection against HB. Contact your doctor, nurse practitioner or public health nurse for information.

A physician or nurse practitioner may recommend that individuals receive HB immune globulin based on an individual's risk assessment of having been exposed to blood and/or body fluids that are known or suspected to be infected with the HB virus through:

- Being poked with an infected needle.
- Being splashed in the mouth, nose or eyes.
- Biting or being bitten by someone with HB.
- Being in contact with personal articles such as razors, toothbrushes and dental floss.
- Using equipment for tattooing, piercings, acupuncture or needles used to inject drugs and other substances (e.g. steroids).

## Who should not get HB immune globulin?

- People who had a life-threatening reaction to a previous dose of an immune globulin product, or any components of the product.
- **Note: Immune globulin can interfere with some vaccines.** Speak with a public health nurse, nurse practitioner or physician if you have been vaccinated for measles, mumps, rubella, chickenpox or shingles within the past 14 days or receive these vaccines in the future.

## What are common reactions to HB immune globulin?

- Temporary pain and tenderness at the injection site.
- Rarely, blot clots may occur after the administration of HB immune globulin.

Acetaminophen (Tylenol, Tempa) may be given for fever or soreness. ASA (Aspirin) must NOT be given to anyone younger than 18 years old because of the risk of Reye syndrome.

It is important to stay in the clinic for 15 minutes after getting HB immune globulin because there is an extremely rare possibility of a life-threatening allergic reaction called anaphylaxis. This may include hives, difficulty breathing, or swelling of the throat, tongue or lips. **If this happens after you leave the clinic, call 911 or the local emergency number.** This reaction can be treated, and occurs in less than one in one million people who get the HB immune globulin.

## Who should you report reactions to?

- Report any adverse or unexpected reactions to your local public health nurse, your doctor, or nurse practitioner as soon as possible.

### Talk to a public health nurse:

- If you have questions or concerns about your or your child's reaction to an immunization.
- If you had to take your child to a doctor, a hospital or to a health centre with a symptom that might be related to immunization.

**For more information contact your local public health office, your physician, nurse practitioner, HealthLine online or by calling 811.**

## What does HB immune globulin contain?

**HepaGam B®** containing human plasma protein, maltose, polysorbate 80, traces of tri-n-butyl phosphate and Triton X-100. Preservative-free.

**HyperHEP B® S/D** contains human hepatitis B hyperimmune globulin, glycine and sodium carbonate. Preservative-free. Prefilled syringes have rubber needle shields and stoppers.

## Mature Minor Consent

It is recommended that parents/guardians discuss consent for treatment with their children. Efforts are first made to get parental/guardian consent for treatment. However, children at least 13 years of age up to and including 17 years of age, who are able to understand the benefits and possible reactions for each product and the risks of not getting treatment, can legally consent to or refuse treatment in Saskatchewan by providing mature minor informed consent to a healthcare provider.

To ensure that a complete immunization record is maintained, every immunization administered to an individual will be documented by Public Health into the electronic provincial immunization registry, known as Panorama. Panorama is a secure electronic system used in Saskatchewan to record and manage immunization records and the health information related to immunization for all Saskatchewan residents. Immunization records may be shared with other health care professionals to provide public health services; assist with diagnosis and treatment; and to control the spread of vaccine preventable diseases. Information about Panorama is detailed in the "Protecting the Privacy of Your Immunization Record" fact sheet.

Reference: Product monographs (HepaGam B® 2016; HyperHEP B® S/D 2012).