

For the Period October 17 to 23, 2017

Harvest has all but wrapped up for producers in the province as 99 per cent of the crop is now combined, according to Saskatchewan Agriculture's weekly Crop Report. There are still some crops such as flax, soybean and sunflower left to be combined.

Harvest weather was favourable for much of the fall, allowing producers to pull off well above-average crop quality. Although field conditions remained dry for the majority of the province, producers had fewer rain delays than in previous years and were able to take most crops off in relatively good condition. There were also limited

reports of diseases such as fusarium head blight impacting crop production this year. The majority of crops are being reported as falling within the top two quality grades.

Crop yields vary greatly throughout the province, depending on seeding date and the amount of moisture received throughout the season. Overall provincial yields are on par with the 10-year average, although some areas in the north are reporting higher than normal yields thanks to timely moisture. Yields in most southern and some central areas of the province were significantly impacted by the extended period of hot and dry conditions this summer. Average provincial yields at this time are reported as 43 bushels per acre for hard red spring wheat, 34 bushels per acre for canola, 18 bushels per acre for soybeans, 63 bushels per acre for barley, 746 lbs per acre for mustard and 1369 lbs per acre for lentils.

Saskatchewan Harvest October 23, 2017 per cent combined	
Winter wheat	100
Fall rye*	100
Spring wheat	100
Durum	100
Oats**	99
Barley	99
Canaryseed	100
Flax	97
Canola	99
Mustard	100
Soybeans	99
Lentils	100
Peas	100
Chickpeas	100
*includes four per cent 'other'	
**includes two per cent 'other'	

One year ago

Harvest progress continued to be stalled across the province due to frequent moisture and saturated fields. Little progress had been made in recent weeks as only 82 per cent of the crop was in the bin. It would be some time before most producers could return to the field; some would not resume combining until the ground froze or until the next spring.

Follow the 2017 Crop Report on
Twitter @SKAgriculture

Saskatchewan Agriculture has a group of 211 volunteer crop reporters from across the province. Thank you for your valued dedication to the crop report. In 2017, there are seven crop reporters reaching their 20 year milestone; three reaching 25 years; three reaching 35 years; and three reaching 40 years.

Congratulations!!

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.saskatchewan.ca/crop-report.



Topsoil and subsoil moisture conditions remain a concern for much of the province. Very little moisture has been received in the last number of weeks and strong winds have further dried fields. The fire risk remains very high in southwestern areas and there have been many reports of grass and stubble fires in recent weeks. Significant moisture will be needed heading into winter to replenish what has been lost throughout the growing season. The majority of producers have indicated that the subsoil is very dry and that seeding conditions next spring will be impacted if moisture is not received. Heading into winter, topsoil moisture on cropland is rated as 40 per cent adequate, 37 per cent short and 23 per cent very short. Hay land and pasture topsoil moisture is rated as 32 per cent adequate, 35 per cent short and 33 per cent very short.

Average hay yields on dry land are reported as 1.1 tons per acre (alfalfa and alfalfa/brome), 0.9 tons per acre (other tame hay), 1.0 ton per acre (wild hay) and 1.6 tons per acre (greenfeed). On irrigated land, the estimated average hay yields are 3.0 tons per acre (alfalfa and other tame hay), 2.0 tons per acre (alfalfa/brome) and 3.5 tons per acre (greenfeed). Hay quality going into winter is rated as 11 per cent excellent, 78 per cent good and 11 per cent fair.

At this time, most livestock producers have indicated that they will have adequate supplies of hay, straw, greenfeed and feed grain heading into winter. However, producers in southern regions are reporting that many areas will have inadequate winter feed supplies and that shortages will be likely. The loss of feed supplies due to recent fires in the southwest has increased concern for producers in those areas.

With the drier than normal field conditions this fall, the number of acres seeded to winter cereals is below average in most areas. Rain was received in much of the province at the end of September, allowing for winter cereal crops to germinate and establish. Thanks to the recent warm and dry weather, producers have been able to complete fall work such as putting down fertilizer, fixing fences, moving cattle, working fields, hauling bales and grain, picking rocks and cleaning up fields. Many producers, particularly in eastern regions, have been able to reclaim acres lost to flooding in previous years.

Saskatchewan Harvest by Crop District October 23, 2017 Per cent combined					
1A	99	3BS	100	7A	100
1B	100	3BN	99	7B	100
2A	100	4A	100	8A	99
2B	99	4B	99	8B	100
3ASE	100	5A	100	9AE	99
3ASW	100	5B	99	9AW	97
3AN	99	6A	100	9B	99
		6B	100		

Provincial Estimated Crop Yields - October 23, 2017								
	Winter wheat	Fall rye	HRSW	Other wheat*	Durum	Oat	Barley	Canary-seed
Southeast	43	41	41	41	34	66	58	949
Southwest	33	27	33	33	33	51	43	873
East Central	50	45	44	47	46	86	67	1,400
West Central	47	34	43	45	45	71	62	1,179
Northeast	65	60	49	54	54	107	71	1,292
Northwest	59	54	47	53	N/A	90	65	N/A
Provincial	43	38	43	46	36	89	63	1,123
10 yr. prov. avg (2007-2016)	42	37	36	N/A	35	75	57	1,211
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea	
Southeast	19	29	621	16	32	1,387	901	
Southwest	17	25	633	13	25	1,238	1,136	
East Central	23	35	1,150	25	40	1,680	N/A	
West Central	25	35	1,182	23	39	1,455	1,200	
Northeast	29	39	N/A	25	45	1740	N/A	
Northwest	25	38	N/A	N/A	41	1394	N/A	
Provincial	21	34	746	18	33	1,369	1,123	
10 yr. prov. ave (2007-2016)	22	31	1,010	N/A	34	1,292	1,265	

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville and Lake Alma areas

Harvest is wrapped up in the region, although there are a few fields of crops such as flax and sunflower left to be combined.

Crop yields vary within the region depending on how much moisture was received throughout the growing season. While producers in the eastern areas received more timely rains, others in western and southern areas were more greatly impacted by the hot and dry conditions. Crops such as soybean, flax, chickpea and mustard were impacted the greatest and yields are well below average. Crop quality is the best it has been in a number of years for producers in the region. The majority of crops are falling within the top two grades thanks to lack of fall moisture and limited disease issues from ergot and fusarium head blight.

Little to no rain was reported last week, allowing producers to complete fall work and get ready for winter. The Alida area has reported the most precipitation (258 mm) in the region since April 1.

Topsoil and subsoil moisture conditions remain a concern for much of the region. With little moisture received in recent weeks coupled with frequent strong winds, topsoil moisture has greatly worsened in the region. Many producers have indicated that the subsoil moisture is also very dry and will need significant amounts of moisture before next spring to replenish what has been lost; seeding conditions will be impacted next spring if field conditions remain dry. Heading into winter, topsoil moisture on cropland is rated as 18 per cent adequate, 38 per cent short and 44 per cent very short. Hay land and pasture topsoil moisture is rated as 14 per cent adequate, 34 per cent short and 52 per cent very short.

Average hay yields on dry land are reported as (in tons per acre): alfalfa and alfalfa/brome 0.9; other tame hay 0.7, wild hay 1.3; and greenfeed 1.4. At this time, most livestock producers have indicated that they will have adequate supplies of hay, straw, greenfeed and feed grain heading into winter. However, some producers in more southern and western areas of the region have indicated that they have inadequate feed supplies and shortages are likely.

With the drier than normal field conditions this fall, the number of acres seeded to winter cereals is well below average in most areas. Although rain was finally received in late September, many producers did not seed winter cereals as fields were still too dry and there were concerns of crops germinating and establishing properly prior to winter. Farmers are busy cleaning up fields, putting machinery away, hauling bales and grain, drying grain, putting down fertilizer, working fields and picking rocks.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Harvest operations are complete in the region, although there are a few fields of flax and sunflower left to be combined.

The lack of moisture and hot temperatures negatively impacted crop production in the region. Crop yields vary greatly from area to area, depending on seeding date and how much moisture was received throughout the growing season. Crops such as canola, mustard, soybeans, field peas and canary seed were the most impacted and yields are well below normal for the region. Crop quality is the best it has been in a number of years, mainly due to lack of fall moisture and limited issues with diseases such as fusarium head blight and root rot. The majority of crops are falling within the top two grades.

Little to no rain was reported last week, allowing producers to complete fall work and get ready for winter. The Moose Jaw area has reported the most precipitation (236 mm) in the region since April 1.

Topsoil and subsoil moisture conditions remain a concern for much of the region. With little moisture received in the last number of weeks coupled with frequent strong winds, topsoil moisture has greatly worsened in the region. The fire risk remains high in much of the region and grass and stubble fires continue to be reported. Subsoil moisture conditions are also very dry and fields will need significant amounts of moisture before next spring. Seeding conditions will be impacted if moisture is not received in the coming weeks. Heading into winter, topsoil moisture on cropland is rated as 17 per cent adequate, 49 per cent short and 34 per cent very short. Hay land and pasture topsoil moisture is rated as nine per cent adequate, 37 per cent short and 54 per cent very short.

Average hay yields on dry land are reported as (in tons per acre): alfalfa, alfalfa/brome and wild hay 0.8; other tame hay 0.7; and greenfeed 1.3. On irrigated land, reported hay yields are 1.3 tons per acre (alfalfa/brome).

At this time, most livestock producers in the region have indicated that they will have adequate supplies of hay, straw, greenfeed and feed grain heading into winter. However, many producers in more southern and western areas of the region have indicated that feed shortages are imminent. Concerns are increasing for producers in the regions as recent fires have destroyed feed supplies as well as grazing land.

The number of acres seeded to winter cereals is well below average in most areas. Although rain was finally received in late September, many producers did not seed winter cereals as fields were still too dry and there were concerns of crops germinating and establishing properly prior to winter.

Farmers are busy working fields, putting machinery away, hauling bales and grain, putting down fertilizer and fixing fences.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Producers have essentially wrapped up harvest operations in the region. There are a few fields of crops such as flax and canola left to be combined. Fall work continues and many producers in the area have been able to reclaim acres lost to flooding in previous years.

Crop yields vary throughout the region but overall are about average to slightly above average thanks to timely rain and good soil moisture during the growing season. Although most crops are about average, crops such as lentils, field peas and durum yielded better than normal for many producers. Crop quality is the best it has been in a number of years, mainly due to lack of fall moisture and limited issues with diseases such as fusarium head blight. The majority of crops are falling within the top two grades.

Small amounts of rainfall were received in the region, although it wasn't enough to keep producers out of the field. The Kelvington area has reported the most precipitation (364 mm) in the region since April 1.

Topsoil moisture conditions remain a concern for much of the region and conditions have worsened in recent weeks. Heading into winter, topsoil moisture on cropland is rated as 58 per cent adequate, 30 per cent short and 12 per cent very short. Hay land and pasture topsoil moisture is rated as 46 per cent adequate, 38 per cent short and 16 per cent very short. Like much of the province, significant amounts of moisture will be needed before next spring in order to replenish what has been lost from the hot and dry conditions.

Average hay yields on dry land are reported as (in tons per acre): alfalfa and alfalfa/brome 1.1; other tame hay and wild hay 1.0; and greenfeed 1.7. At this time, most livestock producers have indicated that they will have adequate amounts of hay, straw, greenfeed and feed grain heading into winter. However, there are some areas that may be short of hay.

The number of acres seeded to winter cereals is below average in most areas. Although rain was finally received in late September, many producers did not seed winter cereals as fields were still too dry and there were concerns of crops germinating and establishing properly prior to winter.

Farmers are busy cleaning up fields, hauling bales and grain, putting fertilizer down, picking rocks and moving cattle.

West-Central Saskatchewan:

- Crop District 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B - Kerrobert, Macklin, Wilkie and Biggar areas

Harvest is all but wrapped up in the west-central region. However, there are a few fields such as canola and flax left to be combined. Fall work continues and many producers in the area have been able to reclaim acres lost to flooding in previous years.

Crop yields vary greatly throughout the region but overall are about average, thanks to timely rain during the growing season. Crops such as lentils, field peas, soybeans and canary seed are slightly above average for the region. Crop quality is the best it has been in a number of years, mainly due to lack of fall moisture and limited issues with diseases such as fusarium head blight. The majority of crops are falling within the top two grades.

Small amounts of rainfall were received in the region, although it wasn't enough to keep producers out of the field. The Cando area has reported the most precipitation (314 mm) in the region since April 1.

Topsoil and subsoil moisture conditions have worsened in the region with the recent strong winds and lack of moisture. Significant amounts of moisture will be needed to replenish what has been lost this summer; seeding conditions will likely be impacted for next spring if fields remain dry. The fire risk remains high in many areas of the region and grass and stubble fires continue to be reported. Heading into winter, topsoil moisture on cropland is rated as 35 per cent adequate, 49 per cent short and 16 per cent very short. Hay land and pasture topsoil moisture is rated as 30 per cent adequate, 50 per cent short and 20 per cent very short.

Average hay yields on dry land are reported as (in tons per acre): alfalfa and alfalfa/brome 0.9; other tame hay and wild hay 0.67; and greenfeed 1.5. On irrigated land, average hay yields are reported as: alfalfa and other tame hay 3.0, alfalfa/brome 3.3; and greenfeed 3.5.

At this time, most livestock producers have indicated that they will have adequate amounts of hay, straw, greenfeed and feed grain heading into winter. However, there are some areas reporting inadequate winter feed supplies and shortages may be likely.

The number of acres seeded to winter cereals is slightly below average in most areas. Although rain was finally received in late September, many producers did not seed winter cereals as fields were still too dry and there were concerns of crops germinating and establishing properly prior to winter.

Farmers are busy working fields, hauling bales and grain, moving cattle, cleaning up fields, fixing fences and putting down fertilizer.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Harvest is virtually complete in the region thanks to a good stretch of warm and relatively dry weather. There are a few fields of crops such as canola, oats and flax left to be combined. Fall work continues and many producers in the area have been able to reclaim acres lost to flooding in previous years.

Crop yields vary throughout the region but overall are well above average for the majority of crops. Timely rain and good soil moisture allowed for crops such as canola, oats, field peas and flax to yield much better than normal for the region. Crop quality is the best it has been in a number of years, mainly due to lack of fall moisture and limited issues with diseases such as fusarium head blight. The majority of crops are falling within the top two grades.

Varying amounts of rainfall were received last week, ranging from trace amounts to 23 mm in the Melfort area. The Nipawin area has reported the most precipitation (595 mm) in the region since April 1.

Topsoil moisture conditions are in relatively good shape heading into winter. The recent rainfall has helped to replenish dry areas, although additional moisture will be needed prior to next spring. Topsoil moisture conditions on cropland are rated as two per cent surplus, 74 per cent adequate, 21 per cent short and three per cent very short. Hay land and pasture topsoil moisture is rated as three per cent surplus, 64 per cent adequate, 31 per cent short and two per cent very short. Crop District 9AE is reporting that 10 per cent of both the cropland and hay land and pasture have surplus topsoil moisture at this time.

Average hay yields on dry land are reported as (in tons per acre): alfalfa 1.9; alfalfa/brome 2.0; other tame hay 1.7; wild hay 1.4; and greenfeed 2.7. At this time, the vast majority of livestock producers have indicated that they will have adequate amounts of hay, straw, greenfeed and feed grain heading into winter.

The number of acres seeded to winter cereals is above average in most areas. Much of the region had adequate topsoil moisture at seeding time and producers were able to take advantage of the quicker-than-normal harvest progress this year.

Farmers are busy finishing up harvest, cleaning up fields, fixing fences, hauling bales and grain, putting fertilizer down, picking rocks and moving cattle.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Ninety-eight per cent of the crop is now combined in the region with the rest of the crop expected to be in the bin within the next week or so. There are some fields of canola, oats and barley left to be combined.

Crop yields vary greatly but overall are average to well above average thanks to timely rain during the growing season. Crops such as spring wheat, flax, canola and field peas are well above average for the region. Crop quality is the best it has been in a number of years, mainly due to lack of fall moisture and limited issues with diseases such as ergot and fusarium head blight. The majority of crops are falling within the top two grades.

Varying amounts of rainfall were received last week, ranging from small amounts to 25 mm in the Neilburg area. The Meadow Lake area has reported the most precipitation (509 mm) in the region since April 1.

Topsoil moisture conditions are in relatively good shape heading into winter. The recent rainfall has helped to replenish dry areas; however, additional moisture will be needed prior to next spring. Topsoil moisture conditions on cropland are rated as 68 per cent adequate, 25 per cent short and seven per cent very short. Hay land and pasture topsoil moisture is rated as 68 per cent adequate, 24 per cent short and eight per cent very short.

Average hay yields on dry land are reported as (in tons per acre): alfalfa 1.5; alfalfa/brome 1.4; other tame hay 1.1; wild hay 0.78; and greenfeed 2.1. At this time, the majority of livestock producers have indicated that they will have adequate amounts of hay, straw, greenfeed and feed grain heading into winter. However, there are reports in some areas of inadequate hay supplies.

The number of acres seeded to winter cereals is above average in most areas. Much of the region had adequate topsoil moisture at seeding time and producers were able to take advantage of the quicker-than-normal harvest progress this year.

Farmers are busy finishing up harvest, putting machinery away, fixing fences, hauling bales and grain, putting fertilizer down and moving cattle.

Saskatchewan Harvest Progress - October 23, 2017

*Other - crop that will not be harvested due to weather, insect or disease damage or will be greenfeed

Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	0	100	
Fall Rye	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	0	0	97	3
southwest	0	0	0	95	5
east central	0	0	0	100	0
west central	0	0	0	100	0
northeast	0	0	0	100	0
northwest	0	0	0	100	0
provincial	0	0	0	96	4
Spring Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	0	100	
Durum	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	N/A	N/A	N/A	N/A	
provincial	0	0	0	100	
Barley	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	1	99	
northeast	0	0	0	100	
northwest	0	0	1	99	
provincial	0	0	1	99	
Oats	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	0	0	98	2
southwest	0	0	0	97	3
east central	0	0	0	99	1
west central	0	1	0	91	8
northeast	0	0	0	99	1
northwest	0	1	0	98	1
provincial	0	1	0	97	2
Canaryseed	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	N/A	N/A	N/A	N/A	
provincial	0	0	0	100	

Flax	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	4	96	
southwest	0	0	4	96	
east central	0	0	0	100	
west central	0	0	2	98	
northeast	0	0	1	99	
northwest	0	0	0	100	
provincial	0	0	3	97	
Canola	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	1	0	99	
west central	0	0	0	100	
northeast	0	1	0	99	
northwest	0	1	0	99	
provincial	0	1	0	99	
Mustard	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	0	0	0	100	
Soybeans	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	1	99	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	1	99	
northeast	0	0	1	99	
northwest	0	0	0	100	
provincial	0	0	1	99	
Field Peas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	0	100	
Lentils	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	0	100	
Chickpeas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	N/A	N/A	N/A	N/A	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	0	0	0	100	

2017 Crop Grades - October 23, 2017

*10 year average is calculated from 2007 to 2016

Winter Wheat				
	1CW	2 CW	3CW	CW feed
2007	63	33	0	4
2008	60	33	0	7
2009	57	36	0	7
2010	28	47	0	25
2011	57	26	0	17
2012	42	31	23	4
2013	42	45	10	3
2014	3	38	44	15
2015	36	45	17	2
2016	33	37	20	10
10 yr avg	42	37	11	9
2017	76	19	5	0

Oats				
	1CW	2CW	3CW	4CW
2007	22	42	26	10
2008	30	54	14	2
2009	27	53	16	4
2010	9	39	36	16
2011	31	48	16	5
2012	20	55	21	4
2013	36	54	9	1
2014	10	62	23	5
2015	19	51	23	7
2016	13	59	18	10
10 yr avg	22	52	20	6
2017	37	57	5	1

Spring Wheat				
	1CW	2 CW	3CW	CW feed
2007	36	39	19	6
2008	50	37	10	3
2009	65	24	8	3
2010	7	29	36	28
2011	54	32	10	4
2012	35	42	16	7
2013	57	32	9	2
2014	9	42	29	20
2015	26	41	23	10
2016	10	42	28	20
10 yr avg	35	36	19	10
2017	77	20	3	0

Rye				
	1CW	2 CW	3CW	sample
2007	67	28	5	0
2008	69	28	3	0
2009	68	23	9	0
2010	29	45	22	4
2011	62	29	9	0
2012	54	38	7	1
2013	53	42	4	1
2014	10	72	12	6
2015	40	53	6	1
2016	65	27	5	3
10 yr avg	52	39	8	2
2017	88	9	3	0

Durum				
	1CW	2 CW	3CW	other (4&5)
2007	46	38	13	3
2008	35	39	19	7
2009	62	26	10	2
2010	3	20	38	39
2011	44	32	17	7
2012	44	32	18	6
2013	21	34	34	11
2014	2	13	37	48
2015	20	40	25	15
2016	4	14	34	48
10 yr avg	28	29	25	19
2017	72	23	4	1

Flax				
	1CW	2 CW	3CW	sample
2007	89	10	1	0
2008	88	11	1	0
2009	85	12	3	0
2010	61	29	7	3
2011	82	14	1	3
2012	87	12	1	0
2013	91	8	1	0
2014	71	21	7	1
2015	73	23	3	1
2016	64	27	7	2
10 yr avg	79	17	3	1
2017	89	10	1	0

Barley			
	Malt	1CW	2CW & sample
2007	43	42	15
2008	48	41	11
2009	35	53	12
2010	14	44	42
2011	42	46	12
2012	24	51	25
2013	36	53	11
2014	19	51	30
2015	22	56	22
2016	26	42	32
10 yr avg	31	48	21
2017	51	42	7

Canola				
	1CAN	2CAN	3CAN	sample
2007	80	16	3	1
2008	90	9	1	0
2009	85	10	3	2
2010	67	19	10	4
2011	82	13	3	2
2012	79	16	4	1
2013	92	7	1	0
2014	74	20	5	1
2015	80	14	4	2
2016	79	14	5	2
10 yr avg	81	14	4	2
2017	91	8	1	0

Mustard				
	1CAN	2CAN	3CAN	sample
2007	73	25	2	0
2008	83	14	3	0
2009	88	10	2	0
2010	64	23	8	5
2011	82	16	2	0
2012	84	12	3	1
2013	86	13	1	0
2014	56	30	12	2
2015	80	18	2	0
2016	64	29	6	1
10 yr avg	76	19	4	1
2017	87	13	0	0

Field Peas				
	1CAN	2CAN	extra 3 &/or 3 CAN	sample
2007	51	43	5	1
2008	44	47	7	2
2009	48	47	4	1
2010	17	49	26	8
2011	39	53	7	1
2012	29	60	10	1
2013	36	61	3	0
2014	13	68	17	2
2015	36	55	8	1
2016	27	60	11	2
10 yr avg	34	54	10	2
2017	61	36	3	0


Soybeans				
	1 CAN	2CAN	3CAN	4&5CAN
2014	33	41	19	7
2015	39	49	10	2
2016	50	41	8	1
2017	38	59	2	1

*2014 is the first year the Crop Report included soybeans

Lentils				
	1CAN	2CAN	extra 3 &/or 3 CAN	sample
2007	45	44	11	0
2008	40	44	14	2
2009	48	45	6	1
2010	5	27	49	19
2011	39	49	11	1
2012	24	54	21	1
2013	35	54	11	0
2014	5	32	53	10
2015	21	54	24	1
2016	4	38	45	13
10 yr avg	27	44	25	5
2017	52	44	4	0


Chickpea				
	1CW	2 CW	3CW	sample
2007	51	43	5	1
2008	48	42	8	2
2009	51	36	11	2
2010	10	28	19	43
2011	46	36	6	12
2012	44	44	11	1
2013	46	44	10	0
2014	13	47	37	3
2015	72	19	8	1
2016	10	36	41	13
10 yr avg	39	38	16	8
2017	78	22	0	0

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

from October 17 to October 23, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

2017 Final Rainfall Summary

in mm

CD	RM	April	May	June	July	Aug	Sept	Oct 1 to 23	Total Yr Precip
1A	2	17	38	62	24	27	52	2	222
	3	17	34	46	50	32	39	0	218
	32	25	38	64	30	42	59	2	258
	33	35	20	61	45	58	47	1	267
	34	7	34	61	29	32	41	13	217
	61	5	16	92	51	74	64	0	302
	64	13	18	85	12	31	32	12	203
	65	19	17	77	4	74	29	9	229
1B	91	2	26	96	16	35	56	3	234
	94	0	1	121	3	47	38	0	210
	122	9	8	96	25	23	62	2	225
	123	4	24	86	11	35	33	5	198
	124	9	33	100	9	31	26	1	209
	125 A	22	8	85	14	12	13	26	182
	125 B	15	18	124	19	19	26	12	233
	151	9	22	96	36	20	38	26	247
	154 A	5	8	89	5	26	11	0	154
	154 B	10	12	85	N/A	N/A	16	0	123
	155	16	8	83	34	23	4	10	177
2A	67	19	2	37	29	18	30	0	135
	68	20	6	38	10	9	24	13	120
	96	0	4	8	8	N/A	13	1	33
	97	10	3	68	7	15.5	8.5	15	127
2B	127 A	26.5	15.5	80	10	22.5	11.5	12	178
	127 B	19	3	20	0	13	13	21	87
	129	10	9	46	2.5	28.5	12	10.5	119.5
	131 A	30	29	38	1	27	9	27	161
	131 B	30	39	39	10	50	30	0	198
	156 A	22.3	6.6	71.2	15.4	25.2	12.4	18	171.1
	156 B	43	12	64	2	64	26	0	211
	159	15	11	52	0	19	14	12	123
	160 A	7	9	34	15	11	12	22	110
	160 B	15	8	28	0	23	15	25	113
	161	21	13	25	5	16	11	35	145
3ASE	162	10.5	16	34.5	14	33	11	19	155
	191	21	13	25	5	16	11	35	126
	38 A	15	2	43	13	13	27	0	113
	38 B	15	1	46	14	13	26	0	114
	39	10	0	34	9	14	18	11	96
3ASW	10	39	7	10	15	13	30	0	114
	12	38	9	9	22	8	9	15	110
	42	23	14	18	23	10	9	16	113
	43	16	20	28	17	11	10	6	108
	73 A	22	27	14	5	10	10	29	117
	73 B	35	24	30	25.5	22	17	31	184.5
	74	0	0	0	1	7	10	29	47
	102	23.5	15.3	19.2	3.1	35	9.6	19.6	125.3
3AN	103	12	29	6	0	15	20	26	106
	132 A	14.5	9.5	20	9	37	11	34.5	135.5
	132 B	30	15.5	79.5	7	48	16	40	236


3BS	193	8	17	32	7	27	14	65	168
	17	13	7.6	5.3	3.3	4.6	1.8	10.9	46.5
	18	20	7	12	25	0	2	26	92
	75	26	22	33	3	31	5	25	145
	76	25	25	14	2	19	11	26	122
	77	40	6	4	26	17	0	36	129
	78	32	16	20	17	4	3	5	97
	105	26	12	40	0	16	12	35	141
	106	26	18	53	6	21	7	25	156
	107	27	7	29	13	7	0	34	117
3BN	108	36	16	7	12	6	17	17	110
	138 A	19	17	62	14	47	5	63	227
	138 B	21	2	56	11	19	8	52	169
	165	0	63	45	30	24	10	73	244
	166	24	21	28	21	10	2	70	176
	168 A	11	20	50	12	20	4	64	181
	168 B	13	9	54	18	29	0	56	179
	226	12	20	19	N/A	N/A	N/A	N/A	51
	228 A	40	26	43	9	25	5	80	228
	257	26	19.5	43	11	30	3	82	214.5
4A	49	34	15	8	35	1	0	N/A	93
	51	31.2	6.2	17.1	26.2	13	4.6	11.7	110
	79	29	17	18	17	8	2	15	106
	109	50	19	14	12	0	0	0	95
	110	36	11	25	7	0	0	0	79
4B	111	26	20	32	11	10	1	50	150
	139 A	15	19	24	34	8	3	48	151
	139 B	20	23	14	13	0	0	0	70
	169	21	37	56	0	0	5	N/A	119
	231	31	21	44	23	41	5	45	210
5A	183	8	25	72	48	0	38	0	191
	211 A	25	28	58	20	91	52	8	282
	211 B	20	15	9	0	N/A	N/A	N/A	44
	213	27	15	57	37	38	15	6	195
	241	10	20	38	31	33	40	15	187
	243	9	4	32	63	23	22	0	153
	244	2	10	60	77	16	2	8	175
	245 A	29	5	38	54	33	17	16	192
	245 B	35	14	35	23	15	15	21	158
	245 C	32.5	19	35	22.5	8	20	19	156
5B	246 A	30	9	53	14	20	16	28	170
	246 B	25	21	56	29.2	31	16.1	16.2	194.5
	247	24	25	45	19	2	0	23	138
	248	17.5	16	46.5	13	8	17	53	171
	271	28	37	32	52	40	31	21	241
	273	34	14	35	64	46	24	12	229
	277	39	36	42	58	30	35	72	312
	305	39	26	37	81	23	36	32	279
	307	26	15	45	65	28	25	46	250
	308 A	13	14	39	22	15	23	61	187
	308 B	13	26	31	28	18	16	77	209
	331	49	25	39	107	47	40	43	350
	335	0	26	21	65	N/A	N/A	N/A	112
	336	31	14	32	85	19	55	11	249
	337	12	29	82	37	28	32	52	277

6A	338	17	26	75	31	6	35	N/A	196
	366	35	30	69	97	36	34	59	364
	367	28	19	78	86	20	31	55	323
	368	22	25	54	121	26	28	42	318
	190 A	24	15	79	8	13	20	29	188
	190 B	30	16	71	0	19	16	28	180
	190 C	38	15	47	0	10	15	39	164
	190 D	29	18	23	0	2	14	21	107
	219 A	22	18	31	3	6	15	30	125
	219 B	41	17	59	2	12	14	N/A	145
	220	19	26	41	10	14	32	19	161
	221 A	26	27	46	25	16	41	26	207
	221 B	14	21	58	10	N/A	23	N/A	126
	222	19	21	40	3	14	22	62	181
	251	7	23	38	25	11	18	70	192
	252	13	20	44	20	20	25	66	208
	279	14.8	22.2	43.3	10	39	28.5	57.4	215.2
	282	16	35	37	12	18	31	104	253
	312	23	31	33	13	22	36	87	245
	313	13	35	49	39	39	33	24	232
	339	23.8	10	53.4	46.2	25.2	25	40.2	223.8
	340	20	37	54	51	44	35	41	282
	341	17	35	21	28	19	N/A	N/A	120
	343	0	35	32	33	54	29	7	190
6B	223 A	11	38	34	2	19	42	47	193
	223 B	5	10	29	12	18	10	80	164
	284 A	18	62	44	31	10	14	53	232
	284 B	16	64	36	31	10	14	53	224
	284 C	15	32	14	83	5	11	32	192
	285	25	28	28	48	7	3	26	165
	286	25	36	11	58	53	4	51	238
	314	10	48	40	10	34	27	41	210
	344	19	58	38	38	52	24	0	229
	346	21	33	30	28	21	19	21	173
7A	376	42	45	34	38	31	19	16	225
	403	20	65	30	63	22	44	15	259
	287	46	10	43	39	27	2	37	204
	288	46	10	45	28	24	52	26	231
	290 A	38	20	46.6	7.1	41.4	2	56.7	211.8
	290 B	25	10	22	5	7	18	50	137
	292	34	27	68	5	38	24	72	268
	317 B	38	23	44	52	17	13	19	206
	318	68	34	51	48	12	0	14	227
	320 A	21	22	71	39	29	14	37	233
7B	320 B	22	30	76	31	28	15	63	265
	321	27	32	66	25	31	27	59	267
	347	44	36	31	27	26	12	20	196
	350 A	40	48	47	60	48	38	22	303
	350 B	40	31	64	21	16	20	25	217
	351	45	51	35	21	50	29	59	290
	352	31	55	46	13	17	20	65	247
	377	38.5	43.5	32	51	35	16	11	227
	378	62	45	39	90	28	24	26	314
	379	38	43	25	23	41	14	23	207
	381	29	34	11	18	59	17	57	225

8A	382	23	53	60	18	45	42	35	276
	409 A	48	79	32	13	18	17	21	228
	409 B	21	77	33	36	57	15	18	257
	410	30.5	20.32	53.34	58.42	68.58	27.94	17.78	276.88
	395	24	61	50	96	11	24	91	357
	397	28.6	37.2	61	112	12.4	18.6	63	332.8
	426	0	0	0	45	0	0	50	95
	428	27	30	41	40	6	18	46	208
	456	31	73	87	59	18	21	64	353
	457	15	107	47	15	2	22	51	259
8B	486	28	78	105	38	13	29	33	324
	487	75	118	225	91	0	38	41	588
	369	15	24	34	36	18	39	2	168
	370 A	19	34	30	44	19	29	40	215
	370 B	15	22	42	40	12	33	32	196
	371	18	38	37	36	30	39	28	226
	372	30.7	57.7	20.5	26.5	37.8	48.8	28.5	250.5
	400	18	29	40	63	19	37	31	237
	429 A	14	57	38	23	2	28	32	194
	429 B	33	43	74	17	1	33	46	247
9AE	459	20	35	23	33	2	22	30	165
	460	13.3	46.5	46.9	19	6.3	13.7	12.7	158.4
	488	32	58	66	34	10	38	10	248
	491	0	24	87	30	0	30	10	181
	520	34	88	28	24	0	18	27	219
	521	34	88	28	24	0	18	27	219
	406	24	34	27	27	77	23	6	218
	435	43	77	41	42	26	48	0	277
	436	33.5	63.5	21	36	25	28	10	217
	463	31.1	52.5	66	35	22	48	13	267.6
9AW	466	53	71	94	45	30	24	16	333
	467 A	54	59	51	56	52	15	9	296
	467 B	24	61	93	56	45	18	13	310
	467 C	48	69	109	61	79	31	17	414
	493	0	0	0	24	18	27	6	75
	494	41	56	60	73	N/A	39	11	280
	497	0	0	0	16	75	0	20	111
	438	52.5	59.5	32	48	41	2	41	276
	440	43.5	46	83	53.5	76	14.5	48.5	365
	442	40.6	72.7	80.5	72.7	50.4	17.6	55.8	390.3
9B	498 A	50	74	59	90	72	6	24	375
	498 B	26	56	32	86	83	6	15	304
	499	51	59	72	85	134	42	15	458
	501 A	95	68	101	82	39	29	24	438
	501 B	31	60	70	82	72	18	18	351
	501 C	63	86	111	60	59	19	16	414
	502	75	77	52	25	39	2	6	276
	561	88	47	95	130	55	15	20	450
	588 A	74	60	147	77	42	34	16	450
	588 B	83	70	149	102	39	36	30	509
	622	141	48	130	127	98	20	13	577


Cumulative Rainfall

from April 1 to October 23, 2017



Cumulative Rainfall


from April 1 to October 23, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions


October 23, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions


October 23, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Minimum Temperature

from October 17 to October 23, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Maximum Temperature

from October 17 to October 23, 2017

